



Tansley review

The small RNA world of plants

Author for correspondence:
 Pierre Rouzé
 Tel: +32 93313800
 Fax: +32 93313809
 Email: Pierre.rouze@psb.ugent.be

Received: 9 March 2006
 Accepted: 3 May 2006

Eric Bonnet¹, Yves Van de Peer¹ and Pierre Rouzé^{1,2}

¹Department of Plant Systems Biology, Flanders Interuniversity Institute for Biotechnology (VIB), Technologiepark 927, B-9052 Gent, Belgium; ²Laboratoire associé de L'Institut National de la Recherche Agronomique, Ghent University, Technologiepark 927, B-9052 Gent, Belgium

Contents

Summary	451	IV. <i>Trans</i> -acting siRNAs	462
I. Introduction	452	V. Conclusion and perspectives	462
II. Small interfering RNAs	453	Acknowledgements	463
III. MicroRNAs	455	References	463

Summary

Key words: noncoding RNA (ncRNA), microRNA (miRNA), RNA interference, *Arabidopsis*, Dicer, plants.

RNA has many functions in addition to being a simple messenger between the genome and the proteome. Over two decades, several classes of small noncoding RNAs *c.* 21 nucleotides (nt) long have been uncovered in eukaryotic genomes, which appear to play a central role in diverse and fundamental processes. In plants, small RNA-based mechanisms are involved in genome stability, gene expression and defense. Many of the discoveries in this new 'small RNA world' were made by plant biologists. Here, we discuss the three major classes of small RNAs that are found in the plant kingdom, namely small interfering RNAs, microRNAs, and the recently discovered *trans*-acting small interfering RNAs. Recent results shed light on the identification, integration and specialization of the different components (Dicer-like, Argonaute, and others) involved in the biogenesis of the different classes of small RNAs in plants. Owing to the development of better experimental and computational methods, an ever increasing number of small noncoding RNAs are uncovered in different plant genomes. In particular the well-studied microRNAs seem to act as key regulators in several different developmental pathways, with a marked preference for transcription factors as targets. In addition, an increasing amount of data suggest that they also play an important role in other mechanisms, such as response to stress or environmental changes.

New Phytologist (2006) **171**: 451–468

© The Authors (2006). Journal compilation © *New Phytologist* (2006)

doi: 10.1111/j.1469-8137.2006.01806.x

I. Introduction

For many years, noncoding RNA (ncRNA) genes were regarded relics of an RNA-based origin of life (Gilbert, 1986; Gesteland *et al.*, 1999). However, as more and more ncRNAs were uncovered, owing to novel experimental and computational approaches (Olivas *et al.*, 1997; Argaman *et al.*, 2001; Huttenhofer *et al.*, 2001; Rivas *et al.*, 2001; Wassarman *et al.*, 2001), it became clear that many of them showed highly specialized biological roles and were not some kind of 'molecular fossils'. In this 'modern RNA world' vision (Eddy, 2001), many ncRNA are involved in functions requiring sequence-specific recognition of another nucleic acid sequence. Such a task is easily performed with RNA molecules through sequence complementarity, fuelling the early idea that ncRNAs would be well suited as regulatory molecules. Indeed, in 1961, François Jacob and Jacques Monod put forward the hypothesis that regulatory genes could produce RNA molecules that would interact with operators by base pairing (Fig. 1), either at the transcriptional level (model I), or the post-transcriptional level (model II). A similar proposal was made a few years later by Britten & Davidson (1969) to explain eukaryotic gene regulation. These views were quickly abandoned after the discovery that protein complexes were involved in the control of almost every step of gene expression. However, the 'RNA breakthrough' in the beginning of this century (Couzin, 2002), with crucial and pioneering contributions of the field of plant biology, was in a way a revival of some of the early ideas.

In 1990, two groups published the same unexpected experimental result, in which overexpression of a gene coding for a chalcone synthase to produce deep purple petunia flowers gave white flowers instead (Napoli *et al.*, 1990; Smith *et al.*, 1990; van der Krol *et al.*, 1990). At that time, this phenomenon, named 'cosuppression', did not find any plausible explanation. Although in plants part of the mystery was solved during the 1990s (Palauqui *et al.*, 1997; Voinnet & Baulcombe, 1997), the molecular mechanism was first discovered in the worm *Caenorhabditis elegans* by studying RNA interference (RNAi; Guo & Kemphues, 1995; Fire *et al.*, 1998; Montgomery *et al.*, 1998; Elbashir *et al.*, 2002). RNAi is present in a broad spectrum of eukaryotes under different names, such as post transcriptional gene silencing in plants (PTGS; Hamilton & Baulcombe, 1999), and quelling in fungi (Cogoni *et al.*, 1996) and algae (Wu-Scharf *et al.*, 2000). The RNAi pathway is thought to act as an immune system against invading nucleic acids coming from viruses, transposons or transgenes (Plasterk, 2002). The silencing is triggered by the presence of long double-stranded (ds) RNA molecules in the cell. These can be synthetic RNAs, replicating viruses or even the result of the transcription of nuclear genes. The dsRNA molecules are chopped in very small pieces of RNA of *c.* 21 nt, referred to as small interfering RNAs (siRNAs), by a specific enzyme named DICER (Bernstein *et al.*, 2001). These siRNAs will be incorporated in an RNA silencing system (RISC) that will recognize, bind and induce cleavage of perfectly complementary mRNAs (Hamilton &




Fig. 1 Jacob and Monod model of RNA-based regulation of gene expression. The regulatory genes produce RNA molecules that will interact with their target genes at the transcriptional (model I) or at the post-transcriptional (model II) level. Reprinted from Jacob & Monod (1961), with permission from Elsevier.

Baulcombe, 1999; Hammond *et al.*, 2000; Hannon, 2002; Zamore *et al.*, 2002). The core component of the RISC complex is a member of the Argonaute protein family which has RNA-binding ability (Hammond *et al.*, 2000). This fundamental discovery makes the artificial silencing of virtually any gene possible with artificially engineered siRNAs, without even the need to know the complete gene sequence, a technique now routinely and widely used in functional genomics. Medical treatments using RNA interference are beginning to be developed (Soutschek *et al.*, 2004).

Another type of small noncoding RNAs are the microRNAs (miRNAs), of which the first one, *lin-4*, was discovered by Victor Ambros and co-workers (Lee *et al.*, 1993). Experiments showed that mutations in the *lin-4* locus disrupted the developmental timing (i.e. normal temporal progression of developmental events) in *C. elegans*. The authors isolated a 693-nt long DNA fragment by positional cloning that could rescue the phenotype of mutant animals. Ambros and colleagues gradually realized (Lee *et al.*, 2004a; Ruvkun *et al.*, 2004 for an accurate and very lively coverage of events), that they were not dealing with a classical protein coding gene but with a tiny ncRNA gene 22 nt long. It was noticed later that the miRNA *lin-4* had antisense complementarity to the RNA sequence of the transcript of the gene *lin-14*, at several places in the 3' UTR region (Wightman *et al.*, 1993). Many classical aspects of miRNA biogenesis were described in those first papers, such as the processing of the small RNA from longer precursor molecules that could form hairpin-like secondary structures, which is now considered as the hallmark of miRNAs. Lee *et al.* (1993) also anticipated that *lin-4* may represent a class of regulatory genes that encode small RNA antisense products. Nevertheless, this astonishing discovery remained unnoticed for some years, and was considered to be an exotic worm-specific process. No evidence for *lin-4*-like miRNAs was found in other organisms and no similar small ncRNAs were detected in nematodes until, in 2000, the miRNA gene *let-7* was found to also act in the developmental timing of *C. elegans* – more in particular in the transition from the first larval stage to the second (Reinhart *et al.*, 2000; Slack *et al.*, 2000). Interestingly, homologs of the *let-7* gene could be identified in the fly and human genomes (Pasquinelli *et al.*, 2000) and only 1 yr later, dozens of novel miRNA genes were identified in flies, human and worms by three different groups (Lagos-Quintana *et al.*, 2001; Lau *et al.*, 2001; Lee & Ambros, 2001).

In conclusion, different small RNA silencing mechanisms have been observed in animals, plants and fungi and are therefore assumed to have evolved from a unique ancestral pathway. However, plants are somehow unusual in the sense that they have highly diversified small RNA (siRNAs, miRNAs and ta-siRNAs) based pathways where other organisms have evolved (or retained) only one. For example, in animals, all the known examples of natural silencing involve miRNAs only. Finally, the budding yeast has apparently lost even the ancestral pathway (Baulcombe, 2004).

In this review we discuss the three major classes of plant small RNA pathways and their functions: small interfering RNAs, microRNAs and the recently discovered *trans*-acting small interfering RNAs.

II. Small interfering RNAs

Small interfering RNAs are generally defined as small RNAs that silence transcripts from which they originate (Bartel, 2004). They were first described in plants, where it was shown that the silencing of three transgenes involved a small antisense RNA c. 25 nt long complementary to each targeted mRNA (Hamilton & Baulcombe, 1999; Hamilton *et al.*, 2002; Tang *et al.*, 2003). In plants, siRNAs have a variety of functions that can be grouped in at least two broad categories: those that trigger changes in the chromatin state of elements from which they derive and those that derive from and defend against exogenous RNA sequences such as viruses or sense transgene transcripts (for review see Baulcombe, 2004).

1. Small interfering RNA silencing of exogenous dsRNA sequences

The siRNAs and miRNA sequences are targeted to a complex called RNA-induced silencing complex (RISC; Fig. 2). Argonaute (AGO) proteins are a core component of this complex (Hammond *et al.*, 2000; Nykanen *et al.*, 2001; Schwarz *et al.*, 2003; Pham *et al.*, 2004; Tomari *et al.*, 2004). In *Arabidopsis*, AGO1 (Bohmert *et al.*, 1998) is associated with miRNAs, *trans*-acting siRNAs and transgene-derived siRNAs but not with virus-derived siRNAs and siRNAs involved in chromatin silencing (Fagard *et al.*, 2000; Boutet *et al.*, 2003; Vaucheret *et al.*, 2004; Kidner & Martienssen, 2005). Mutants of AGO1 were shown to be hypersensitive to virus infections (Morel *et al.*, 2002). Some transposons were shown to be upregulated in *ago1* mutants (Lippman *et al.*, 2003). It was also shown that at least *in vitro*, AGO1 does not seem to have other partners in RISC and would be solely interacting with small RNAs, unlike what is observed in animals (Baumberger & Baulcombe, 2005).

The RNA-dependent RNA polymerases (RDRs) RDR1 and RDR6 are required in the siRNA pathway that silences viruses and transgenes (Dalmay *et al.*, 2000; Mourrain *et al.*, 2000; Xie *et al.*, 2001). Those proteins turn single-stranded (ss) RNA into dsRNA, with or without a siRNA as a primer (Baulcombe, 2004). As a result of the action of RDRs, a single RNA or primary siRNA molecule can generate many dsRNA, thus amplifying the response (Fig. 2b). Unexpectedly, it was shown that RDR6 might also repress the expression of a miRNA, miR165/166 (Li *et al.*, 2005).

The dsRNA structures are further processed by a member of the Dicer family that generates small RNAs from double-stranded RNA sequences with 2-nt overhangs at the 3' ends (Bernstein *et al.*, 2001). In *Arabidopsis*, four Dicer-like (DCL)




Fig. 2 Small RNA pathways in plants. (a) Plant microRNA (miRNA) biogenesis. MicroRNA genes are transcribed from their own locus by POL-II. The hairpin-like secondary structure is further processed by DICER in several steps to produce miRNA:miRNA* duplexes. The duplexes are then methylated by HEN1, before being exported to the cytoplasm, possibly by HASTY. Here the duplex is unwound and the miRNA is associated with AGO1. This complex, known as RISC, will bind specifically to a target messenger RNA, and guide its cleavage (in most of the cases) or will repress its translation. DCL, Dicer-like; HYL, HYPONASTIC LEAVES. (b) Small interfering RNAs (siRNAs). Long double-stranded RNAs (dsRNAs) from diverse origins (viruses, transposons, transgenes, etc.) are converted into 21 nucleotide (nt) long siRNAs by DICER enzymes. These small RNAs are then loaded into RISC and associated with AGO4 or another Argonaute protein. The complex will then bind to the same messenger RNA from which they originate, and cleave the mRNA, silencing its expression. Small interfering RNAs can also bind to the mRNA and initiate the transformation of single-stranded RNA (ssRNA) into dsRNA, thus amplifying siRNA production. RDR, RNA-dependent RNA polymerase. (c) *trans*-Acting siRNAs. In plants, some miRNAs (miR173 and miR390) cleave a target mRNA expressed from ta-siRNA loci. After cleavage, either the 5'- or the 3'-terminus is converted into dsRNA by RDR enzymes, and then processed into 21 nt siRNAs that guide degradation of target mRNA that is different from the ta-siRNA transcript from which they originated.

proteins are known, each one having a distinct function in different small RNA pathways (Schauer *et al.*, 2002; Xie *et al.*, 2004; Dunoyer *et al.*, 2005; Gascioli *et al.*, 2005; Xie *et al.*, 2005b). DCL1 produces miRNAs, DCL2 produces siRNAs involved in the silencing of at least some viral sequences (Xie *et al.*, 2004) and DCL3 produces siRNAs involved in DNA methylation and heterochromatin formation (Xie *et al.*, 2004). Recent work suggests that DCL4 produces siRNAs triggered by inverted-repeat transgenes in plants (Dunoyer

et al., 2005) and is also associated with the ta-siRNA pathway (Gascioli *et al.*, 2005; Xie *et al.*, 2005b; Yoshikawa *et al.*, 2005). Some results have also shown that a partial functional redundancy amongst the different Dicer-like proteins in *Arabidopsis* is possible (Gascioli *et al.*, 2005; Xie *et al.*, 2005b).

Virus dsRNA sequences are recognized by the RNA silencing machinery, which produces siRNAs that will silent viral genes and prevent the accumulation of the pathogen (for review see Dunoyer & Voinnet, 2005). Virus defense via

siRNA is likely to be an ancient mechanism and therefore viruses have evolved various ways to bypass this barrier (Baulcombe, 2004; Dunoyer & Voinnet, 2005; Simon-Mateo & Garcia, 2006). The mechanism best known involves the production of a protein by the virus that will block the silencing pathway of the host (Voinnet *et al.*, 1999; Mallory *et al.*, 2002; Kasschau *et al.*, 2003; Ye *et al.*, 2003; Chapman *et al.*, 2004; Dunoyer *et al.*, 2004; Lakatos *et al.*, 2004). But other mechanisms might also exist. In plants, the expression of siRNAs from inverted-repeats transgenes mimic the symptoms observed during viroid infections, suggesting RNA silencing of the host genes (Wang *et al.*, 2004b). Conversely, many *Arabidopsis* siRNAs do not show a high degree of similarity to any *Arabidopsis* mRNA. Therefore, one interesting hypothesis is that they could constitute a reservoir of defense molecules because of their complementarity to viral sequences (Dunoyer & Voinnet, 2005). This RNA silencing defense might not be limited to viruses: it was also shown recently that bacterial infection by a virulent *Agrobacterium tumefaciens* triggered a rather complex siRNA-mediated silencing response (Dunoyer *et al.*, 2006).

Another striking feature of siRNA-mediated silencing in plants and in some animals is its systemic nature: the effect of silencing can extend beyond the site of initiation and spread through the organism (for a review see Voinnet, 2005). Recent work suggests that DCL4 is responsible for the production of the 21-nt long siRNAs involved in the cell-to-cell silencing signal (Dunoyer *et al.*, 2005). The movement of siRNAs or miRNAs could be important for the regulation of endogenous genes. For example, it is known that the distribution of miR165/166 in the leaf, where they act as repressors of genes affecting leaf polarity, resembles that of a mobile signal (Emery *et al.*, 2003; Juarez *et al.*, 2004; Kidner & Martienssen, 2004). Many siRNAs and miRNAs were detected in the phloem sap of pumpkin, where a protein has been characterized that binds specifically to small RNAs (Yoo *et al.*, 2004).

Recently, a study revealed that an antisense overlapping gene pair generated two types of siRNAs involved in salt-stress tolerance (Borsani *et al.*, 2005). Those genes are *P5CDH*, a stress-related gene, and *SRO5*, a gene of unknown function. When both transcripts are present, a 24-nt siRNA is formed by a biogenesis pathway dependent on DCL2, RDR6, SGS3 and NRPD1A. The cleavage of the *P5CDH* transcript by the 24-nt siRNA then sets the phase for the generation of 21-nt siRNAs by DCL1 that will further cleave the *P5CDH* transcript. The expression of *SRO5* is induced by salt, a step thus necessary for the initial siRNA formation. This elegant study shows that endogenous siRNAs (dubbed nat-siRNAs), derived from a pair of natural *cis*-antisense transcripts, regulate salt tolerance. Given that overlapping genes are not rare in many eukaryotic genomes, nat-siRNA-based regulation might also occur in many other processes (Borsani *et al.*, 2005).

2. Small interfering RNAs and chromatin state

The link between siRNAs and chromatin modifications was mostly explored in the fission yeast *Saccharomyces pombe* where it appears that siRNA-mediated heterochromatin modification is a general mechanism for regulating gene expression (for a review see Lippman & Martienssen, 2004; Gendrel & Colot, 2005). The existence of a mechanism of *de novo* methylation of genes that can be induced and targeted in a sequence-specific manner was first shown in plants (Wassenegger *et al.*, 1994). Later, a link between locus specific siRNAs and histone modifications (deacetylation) or histone H3 lysine 9-methylation was later shown in plants (Aufsatz *et al.*, 2002; Jackson *et al.*, 2002; Zilberman *et al.*, 2003; Xie *et al.*, 2004).

Experiments have shown that expressed siRNAs were matching transposable element sequences that could form imperfect RNA duplexes (Mette *et al.*, 2002). More recently, high-throughput sequencing of expressed small RNAs in *Arabidopsis* (Lu *et al.* 2005a) showed that many siRNAs are associated with transposons silenced by methylation. Maintaining this silenced state involves a low level of transcription, which is a paradox because silencing inhibits transcription by known DNA-dependent RNA polymerases (Lippman & Martienssen, 2004). However, the recent finding of a new RNA polymerase might help to solve this issue. Two studies have described a new RNA polymerase named POL-IV that directs heterochromatic silencing, although the mechanism is not yet clear (Herr *et al.*, 2005; Kanno *et al.*, 2005; Onodera *et al.*, 2005; Pontier *et al.*, 2005; Vaughn & Martienssen, 2005). The RNA-dependent RNA polymerase RDR2 was shown to be required for heterochromatin formation, as well as the Dicer-like protein DCL3 and the Argonaute protein AGO4 (Zilberman *et al.*, 2003; Xie *et al.*, 2004; Zilberman *et al.*, 2004).

Unexpectedly, similarly to siRNAs, miRNAs might also contribute to DNA methylation in *Arabidopsis*. There is experimental evidence that mir165/166, which targets the PHAVOLUTA (PHV)- and PHABULOSA (PHB)-encoding mRNAs, induces methylation of *PHV* and *PHB* genes downstream of the miRNA target sites (Bao *et al.*, 2004). MicroRNA pairing very likely takes place with the nascent but already spliced transcript, implying that miRNAs may also be active in the nucleus, at least in plants.

III. MicroRNAs

1. Biogenesis of miRNAs

Plant miRNAs were first identified in early 2002 (Llave *et al.*, 2002a; Park *et al.*, 2002; Reinhart *et al.*, 2002). Like their animal counterparts, they are short sequences *c.* 21 nt long, processed from longer precursor sequences. While animal precursor sequences usually have a length of 70–80 nt, plant miRNA precursor sequences are much more variable and




Fig. 3 Plant microRNA (miRNA) stem-loop secondary structures examples from *Arabidopsis* (ath-miR156a, ath-miR319a and ath-miR162a), *Oryza* (osa-miR162a) and *Populus* (ptc-miR162a). The mature miRNA sequence is shown in bold letters. Although all structures share a stem-loop structure, the shape can vary slightly. Close homologs of the *Arabidopsis* miRNA ath-miR162a have been found in rice (osa-162a) and poplar (ptc-162a). In those cases, the mature miRNA sequence is exactly the same, while the rest of precursor sequence differs.

range from 50 nt to more than 350 nt. First discovered in *Arabidopsis thaliana*, some plant miRNAs were found to be conserved in many plant genomes such as those of *Oryza sativa*, *Zea mays* and those of more ancient vascular plant genera such as ferns or even nonvascular plants such as mosses (Floyd & Bowman, 2004; Axtell & Bartel, 2005).

Among the main differences between plant siRNAs and miRNAs is that the latter are processed from their own loci (Fig. 2a). In plants, primary miRNA transcripts (pri-miRNA) are produced by RNA polymerase II (POL-II) and are capped

and polyadenylated (Aukerman & Sakai, 2003; Xie *et al.*, 2005a). A TATA regulatory binding motif was found in the upstream region of at least some *Arabidopsis* pri-miRNAs (Kurihara & Watanabe, 2004; Xie *et al.*, 2005a). Since pri-miRNAs are polyadenylated, some of them can be found in Expressed Sequences Tags (EST) databases (Jones-Rhoades & Bartel, 2004). The mature miRNA sequence can be found on either the 5' or the 3' strand of the precursor sequence (Reinhart *et al.*, 2002; see, for example, ath-miR156a and ath-miR319a in Fig. 3). One mature miRNA is encoded by one

or more miRNA genes and sequences that only differ by a few nucleotides are usually grouped together to form families (Reinhart *et al.*, 2002; Griffiths-Jones *et al.*, 2006). Transgenic experiments have shown that it was possible to replace the miRNA:miRNA* duplex by an artificial hairpin structure without altering miRNA processing, thus showing that structure is more important than the sequence itself in this process (Parizotto *et al.*, 2004; Vaucheret *et al.*, 2004). Statistical analyses have shown that pre-miRNA secondary structures tend to have free energy values that are significantly different from those of random sequences, contrary to structures of other classes of ncRNAs, such as transfer RNAs or ribosomal RNAs (Bonnet *et al.*, 2004b; Clote *et al.*, 2005). This indicates that miRNA precursor sequences have highly stable secondary structures, a property likely necessary to avoid anticipated degradation and to allow correct processing by Dicer enzymes.

In *Arabidopsis*, miRNA biogenesis in the nucleus is performed in several steps and requires both DCL1 and HYL1 (Fig. 2a; Papp *et al.*, 2003; Kurihara & Watanabe, 2004; Kurihara *et al.*, 2006). Another protein, HEN1, is required for miRNA biogenesis. This enzyme has two dsRNA-binding domains and a nuclear localization signal. It is conserved in fungi and is required for miRNA and siRNA processing (Park *et al.*, 2002; Boutet *et al.*, 2003; Xie *et al.*, 2003). It was shown recently that HEN1 is responsible for the 3' end methylation of *Arabidopsis* miRNA:miRNA* duplexes and that this modification could be essential for their biogenesis and function in the RNA silencing pathway. All known classes of endogenous small RNAs in *Arabidopsis* require HEN1 (Yu *et al.*, 2005).

HYPONASTIC LEAVES 1 (HYL1) is required to process miRNAs and has a nuclear localization signal (Lu & Fedoroff, 2000; Han *et al.*, 2004; Vazquez *et al.*, 2004a). It was shown that HYL1 interacts with DCL1 *in vitro* (Hiraguri *et al.*, 2005). HASTY (HST, Bollman *et al.*, 2003) may be involved in export of miRNA to the cytoplasm, although convincing evidence for it is lacking, but HST-independent nucleocytoplasmic pathways do exist since miRNA export is not totally blocked in *hst* mutants (Park *et al.*, 2005).

2. Identification of plant miRNA genes and their targets

Arabidopsis miRNA genes were first identified using cloning experiments, by which small RNAs (size between 16 nt and 30 nt) are isolated from whole plants and then cloned and sequenced (Llave *et al.*, 2002a; Park *et al.*, 2002; Reinhart *et al.*, 2002; Sunkar & Zhu, 2004; Gustafson *et al.*, 2005). As a consequence, different types of small RNAs such as siRNAs, miRNAs and RNA degradation products are selected with this procedure. In order to select for miRNAs, the secondary structure of the miRNA precursor sequence is checked for compliance with known miRNA features (Ambros *et al.*, 2003). To extract the potential precursor sequence, small RNAs are mapped back on genomic sequences to extract

flanking regions. The secondary structure is then predicted using *ad hoc* software tools, such as the MFOLD or RNAFOLD packages (Zuker & Stiegler, 1981; Hofacker *et al.*, 1994). Using cloning experiments, miRNA sequences were identified in *Arabidopsis* (Reinhart *et al.*, 2002; Sunkar & Zhu, 2004), rice (Sunkar *et al.*, 2005), poplar (Lu *et al.*, 2005b), moss (Arazi *et al.*, 2005) and tobacco (Billoud *et al.*, 2005). However, this kind of approach implies long and tedious bench work, and the detection is restricted to the most abundant molecules. MicroRNAs expressed under special conditions (stress, etc.) or at specific points in time will remain undetected.

Complementary to cloning experiments are computational approaches for the prediction of miRNAs. Here, the main problem is in discriminating between real miRNAs and so-called false positives. Many algorithms start with predicting all possible hairpin structures for a given genomic sequence (Lim *et al.*, 2003). However, the number of such structures is usually very high, with many false positives, for example, because of the repeats present in most genomic sequences. Therefore, most approaches add to this first step several filters based on the properties of experimentally documented miRNAs. One of the properties used by most algorithms is evolutionary conservation. It has indeed been shown that many miRNAs are conserved in different organisms (Bartel, 2004). In plants, the conservation is, in most cases, limited to the mature miRNA (*c.* 21 nt) while the rest of the precursor sequence is far less well conserved (Fig. 3). For example, even between distantly related plants such as *Arabidopsis* and rice that have diverged more than 130 millions years ago (Friis *et al.*, 2004), the sequence of miR162 is completely conserved (Fig. 3; Reinhart *et al.*, 2002). Parameters describing the secondary structure such as free energy, the number of paired residues within the miRNA, or the number and the size of bulges are used to select valid structures, with cut-off values based on experimentally proven miRNAs. Compositional characteristics such as GC content or the low-complexity content of the miRNA sequences can also help to get rid of irrelevant repeat sequences (Bonnet *et al.*, 2004a; Jones-Rhoades & Bartel, 2004; Wang *et al.*, 2004a; Adai *et al.*, 2005).

Initially, target prediction for plant miRNAs was quite straightforward because it was assumed that most of them match their targets with near-perfect complementarity (Rhoades *et al.*, 2002). In that case, the search for transcript targets is done for a sequence pattern complementary to a given miRNA sequence, allowing few mismatches (usually two or three). For example, Jones-Rhoades & Bartel (2004) used a score system taking into account mismatches, gaps and G:U base pairs. However, more recent experimental work on miRNA targets in *Arabidopsis* showed that some miRNA match their targets with even less complementarity and that the binding pattern is not random (Mallory *et al.*, 2004b; Allen *et al.*, 2005; Schwab *et al.*, 2005). As in animals, there are fewer mismatches in the 5' part of miRNA:mRNA while

the free energy of the duplex is defined by a maximum value. Many miRNA computational prediction tools integrate target detection to support the result of miRNA prediction. In some cases, the target prediction is restricted to one genome (Bonnet *et al.*, 2004a; Wang *et al.*, 2004a; Adai *et al.*, 2005). In other cases, the prediction is further constrained through conservation of targets in several genomes (Jones-Rhoades & Bartel, 2004). So far, computational pipelines have been applied successfully for the discovery of new conserved miRNAs using the complete genomes of *A. thaliana* and *O. sativa* (Bonnet *et al.*, 2004a; Jones-Rhoades *et al.*, 2004; Wang *et al.*, 2004a; Adai *et al.*, 2005).

Many of the predicted targets have been verified experimentally. A modified version of the 5' RACE is usually applied to look for the product of degradation consecutive to the cleavage of the targeted miRNA (Llave *et al.*, 2002b; Jones-Rhoades *et al.*, 2004; Lu *et al.*, 2005b). Some studies remapped known miRNAs on newly available plant genomic sequences such as *Sorghum bicolor* (Bedell *et al.* 2005). The availability of new plant genomes, like the poplar (<http://genome.jgi-psf.org/Poptr1/Poptr1.home.html>; Tuskan *et al.*, 2004), should provide new material to which new computational searches for conserved miRNAs can be applied.

Most of the experimentally documented miRNA sequences from plant, animal and viral genomes are deposited in the miRNA registry (Griffiths-Jones *et al.*, 2006). Release 7.1 (October 2005), contains 731 plant miRNAs genes, representing eight plant genomes (Table 1). The largest number of miRNA families is found in *Arabidopsis* and rice, with 46 families encoded, respectively, by 117 and 178 miRNA genes. Next is poplar with 33 miRNA families encoded by 213 miRNA genes. Those three genomes are the only complete plant genomes currently available, so it is no surprise that they have the largest collection of miRNA genes. A typical feature for plant miRNAs is that they can be divided in two broad categories: miRNAs that are conserved in different plant genomes and miRNAs that are specific to one organism (Reinhart *et al.*, 2002; Jones-Rhoades & Bartel, 2004; Sunkar & Zhu, 2004; Sunkar *et al.*, 2005). The existence of organism-specific miRNAs seems to be specific for plants, although a recent study suggested the existence of a pool of primate-specific miRNAs (Bentwich *et al.*, 2005).

3. MicroRNA function

Most of the plant miRNAs play a role in developmental processes, and a majority of their targets are transcription factors (see for review Jones-Rhoades *et al.*, 2006). It now appears that at least some of the miRNAs are involved in other processes as well, such as response to environmental conditions (Jones-Rhoades & Bartel, 2004; Fujii *et al.*, 2005; Lu *et al.*, 2005b; Sunkar *et al.*, 2005; Chiou *et al.*, 2005). Plant miRNAs seem to be much more specific than their animal counterparts (Schwab *et al.*, 2005). It is estimated that in humans, miRNAs

Table 1 Plant microRNA (miRNA) distribution

	miRNA families	miRNA genes
<i>Arabidopsis thaliana</i>	46	117
<i>Oryza sativa</i>	46	178
<i>Populus trichocarpa</i>	33	213
<i>Zea mays</i>	18	97
<i>Sorghum bicolor</i>	16	72
<i>Glycine max</i>	11	22
<i>Medicago truncatula</i>	10	16
<i>Saccharum officinarum</i>	6	16

Data from the miRNA registry release 7.1 (Griffiths-Jones *et al.*, 2006). Genomes that have been fully sequenced are indicated in bold type.

could regulate up to one-third of the protein coding genes (Lim *et al.*, 2005). Contrary to their animal counterparts, plant miRNA targets are more often located in coding sequences and only occasionally in UTR regions (Bartel, 2004). There are several ways to decipher the biological role of a given miRNA. The simplest way consists of finding the target gene(s) of the miRNA, experimentally and/or through *in silico* search. More sophisticated approaches involve the use of mutants, knock-outs or overexpression of the miRNA precursor genes.

Experimental validation for predicted targets was facilitated by the fact that most of the plant miRNA targets are regulated by cleavage, allowing detecting products of degradation with *ad-hoc* experiments (modified 5' RACE, see Llave *et al.*, 2002b; Jones-Rhoades & Bartel, 2004; Lu *et al.*, 2005b). Several groups went a step further and built artificial constructs in order to overexpress a given miRNA or to make predicted targets resistant to miRNA matching by the introduction of mutations. Here we try to summarize those results and to group them according to the biological process in which the miRNA is involved (see also Table 2).

Leaf, floral and shoot development An overexpression of miR172 in *Arabidopsis* was demonstrated to cause early flowering and defects in floral identity such as absence of petals and transformation of sepals into carpels (Aukerman & Sakai, 2003). The predicted targets of miR172 are members of the APETALA2 transcription factors, including AP2 itself, but also TARGET OF EAT (TOE1 and TOE2) or GLOSSY15 (GL15). Loss of function analyses for these genes indicate that they normally act as floral repressors. The downregulation of AP2-like genes by miR172 during the early stages of development relieves floral repression and promotes flowering. Surprisingly, in *Arabidopsis* miR172 appears to downregulate its targets through a mechanism of translational repression rather than cleavage. Indeed, products corresponding to the cleavage of AP2-like targets by miR172 were found by different groups, but they may represent a small fraction of the total AP2 transcript population

Table 2 Functions of conserved plant microRNAs (miRNAs) for which computational prediction and experimental evidence have been published

Function	miRNA families	Target gene(s)	miRNA conservation	References
Floral timing and leaf development	156	SPL transcription factors	A G O P Sb So	26
Floral development	171	SCL-like transcription factors	A O P S	24, 25
Floral development and vegetative phase change	172	AP2-like transcription factors (AP2, TOE1, TOE2, GL15)	A G O P Sb Z	1, 6, 20, 26
Expression of Auxin response genes, developmental defects	160	160: Auxin response factors (ARF10, ARF16, ARF17)	160: A G M O P Sb Z	8, 19, 21, 23, 26, 27
	164	164: NAC1	164: A O P Sb Z	
	167	167: ARF6, ARF8	167: A G O P So Sb Z	
	393	393: TIR1	393: A M O P Sb Z	
	390	390: ARF3, ARF4 via TAS3 (ta-siRNA)	390: A O P	
Organ separation and number	164	CUC transcription factors	A O P Sb Z	11, 12, 18
Organ polarity, vascular and meristem development	165/166	HD-ZIP III transcription factors (PHB, PHV, leaf 1)	A G M O P Sb Z	2, 7, 9, 10, 13, 14, 16, 17
Floral and leaf patterning	159/319	319: TCP transcription factors 159: MYB transcription factors	A G M O P S S Z	3, 5, 22
Regulation of the miRNA pathway	162	162: DCL1	162: A M O P Z	4, 15, 27
	168	168: AGO1	168: A G O P So Sb Z	
	403	403: AGO2	403: A P	
Sulfate assimilation	395	ATP sulfurylases (APS1, APS3, APS4)	A O P S Z	8
Lignin formation?	397	Laccases	397: A O P	8, 25, 32
	408			
Oxidative stress	398	Copper superoxide dismutases (CSD1, CSD2)	A G O P	8
Phosphate homeostasis	399	Phosphate transporter, E2 ubiquitin-conjugating enzyme	A M O P Sb Z	8, 36, 37
Unknown	161	161: PPR	161: A	8, 27, 28, 29, 30, 31, 33, 34, 35
	163	163: SAM-dependent methyl transferase	163: A	
	173	173: PPR via TAS1 & TAS2 (ta-siRNA)	173: A	
	390	390: Receptor-like kinase	390: A O P	
	394	394: F-box	394: A O P Sb Z	
	396	396: Growth response factors (GRF1, GRF2, GRF3, GRF7, GRF8, GRF9), rhodanese	396: A G O P So Sb Z	

The miRNA conservation in plant species is given according to the miRNA registry data, release 7.1 (Griffiths-Jones *et al.*, 2006).

Species abbreviations: A, *Arabidopsis thaliana*; G, *Glycine max*; M, *Medicago truncatula*; O, *Oryza sativa*; P, *Populus trichocarpa*; So, *Saccharum officinarum*; Sb, *Sorghum bicolor*; Z, *Zea mays*.
References: 1, Aukerman and Sakai (2003); 2, Emery *et al.* (2003); 3, Palatnik *et al.* (2003); 4, Xie *et al.* (2003); 5, Achard (2004); 6, Chen (2004); 7, Floyd and Bowman (2004); 8, Jones-Rhoades & Bartel (2004); 9, Juarez *et al.* (2004); 10, Kim *et al.* (2004); 11, Laufs *et al.* (2004); 12, Mallory *et al.* (2004a); 13, Mallory *et al.* (2004b); 14, McHale and Koning (2004); 15, Vaucheret *et al.* (2004); 16, Williams *et al.* (2005b); 17, Zhong and Ye (2004); 18, Baker *et al.* (2005); 19, Guo *et al.* (2005); 20, Lauter *et al.* (2005); 21, Mallory *et al.* (2005); 22, Millar and Gubler (2005); 23, Wang *et al.* (2005); 24, Llave *et al.* (2002b); 25, Schwab *et al.* (2005); 26, Kasschau *et al.* (2003); 27, Allen *et al.* (2005); 28, Vazquez *et al.* (2004); 29, Rhoades *et al.* (2002); 30, Sunkar *et al.* (2004); 31, Sunkar & Zhu (2005); 32, Lu *et al.* (2005b); 33, Peragine *et al.* (2004); 34, Yoshikawa *et al.* (2005); 35, Park *et al.* (2002); 36, Chiou *et al.* (2006); 37, Fujii *et al.* (2005).

(Aukerman & Sakai, 2003; Kasschau *et al.*, 2003; Chen, 2004; Schwab *et al.*, 2005). Analysis of transgenic maize lines overexpressing GL15 showed that this gene controls the transition from juvenile to adult leaves (Lauter *et al.*, 2005). This transition is the result of opposite effects of GL15 and miR172, the latter promoting the transition to the adult phase by downregulating GL15. Data suggest that this could be a general mechanism for the regulation of vegetative phase change in higher plants (Lauter *et al.*, 2005).

The miRNA miR171 is perfectly complementary to three members of the *SCARECROW*-like family of transcription factors (*SCL6-II*, *SCL6-III* and *SCL6-IV*). This gene family controls a wide range of developmental processes, including radial patterning in roots and hormone signaling. The cleavage of *SCL6-III* and *SCL6-IV* by miR171 was shown by 5' RACE experiments. The fact that *SCL6-III* and *SCL6-IV* are predominantly found in inflorescence tissues, just like miR171, might suggest a role for this miRNA in flowering processes but this hypothesis needs to be confirmed by specific experiments (Llave *et al.*, 2002b).

The *SPL* genes encode a class of plant-specific transcription factors (SQUAMOSA PROMOTER BINDING PROTEIN LIKE) that were predicted to be the targets of miR156 (Rhoades *et al.*, 2002). The overexpression of this miRNA has been shown to cause a moderate delay in flowering and a faster initiation of rosette leaves compared with wild-type. A severe decrease of apical dominance is also observed and the first flowers tend to arise from side shoots. The combination of those traits leads to a phenotype with a substantial increase (up to 10 times) in total leaf number on the side of the shoots (Schwab *et al.*, 2005).

The transcription factor genes *cup-shaped cotyledons* (*CUC*) are predicted to be the targets of the miR164 family. Expression of miR164-resistant versions of *CUC1* caused alterations in *Arabidopsis* embryonic, vegetative and floral development, affecting cotyledon orientation, rosette leaves shape, petals and sepals number (Mallory *et al.*, 2004a). Overexpression of miR164 reproduced the phenotype of *cuc1 cuc2* double mutants by downregulating the levels of *CUC1* and *CUC2* but not *CUC3* mRNAs (Laufs *et al.*, 2004; Mallory *et al.*, 2004a). Disruption of the regulation of *CUC2* by miR164 caused enlarged sepal boundary domains, indicating that miR164 regulation constrains the expansion of the boundaries by degrading *CUC1* and *CUC2* mRNAs (Laufs *et al.*, 2004). Analysis of the mutant *early extra petals 1* (*eep1*) that was found to encode for an extra member of the miR164 family (miR164c) revealed that this miRNA controls petal numbers by regulating *CUC1* and *CUC2* transcript accumulation (Baker *et al.*, 2005).

MiR319 (also known as miR-JAW) was identified through a genetic screen and guides the cleavage of several *TCP* transcription factor genes controlling leaf development (Palatnik *et al.*, 2003). Mutants for the miR319 locus exhibited crinkled leaves, as well as an overexpression of miR319. Constitutive

expression of *TCP2* or *TCP4* partly rescued the miR319 mutant, with leaves less affected but still different from the wild type. MiR159 is a close homolog of miR319 that differs by only three residues and guides the cleavage of two transcripts encoding MYB transcription factors (*MYB33* and *MYB65*). *Arabidopsis* plants transformed with cleavage resistant *MYB33* exhibit pleiotropic developmental defects (Palatnik *et al.*, 2003; Millar & Gubler, 2005). A feedback regulation of *MYB* genes on miR159 levels was revealed in *Arabidopsis*, as part of the gibberellin–DELLA proteins controlling flower development (Achard *et al.*, 2004).

Leaf polarity, vascular and meristem development Members of the class III HD-ZIP transcription factor gene families *PHABULOSA* (*PHB*) and *PHAVULOTA* (*PHV*) govern vascular pattern and leaf polarity. Both gene families have complementary sites for miR165/166. Regulation by those miRNAs was indeed shown to be necessary for a proper organ axis specification, vascular development and meristem function (Emery *et al.*, 2003; Mallory *et al.*, 2004b; McHale & Koning, 2004; Zhong & Ye, 2004; Kim *et al.*, 2005; Williams *et al.*, 2005b). In maize, it was demonstrated that miR166 is a conserved polarizing signal whose expression pattern spatially defines the expression of the HD-ZIP III member *ROLLED LEAF 1*, determining the abaxial (upper) and adaxial (lower) asymmetry of the leaf (Juarez *et al.*, 2004). Moreover, the cleavage of HD-ZIP III genes by miR165/166 was found to be extremely well conserved amongst vascular plants, including ferns and mosses (Floyd & Bowman, 2004).

Auxin response Auxin is a phytohormone implicated in virtually every aspect of plant growth and development. Most of its effects are mediated by *auxin transcription factor* (*ARF*) genes. In *Arabidopsis*, this family consists of 23 genes. Among those, three were predicted to be targets of miR160, namely *ARF10*, *ARF16* and *ARF17* (Rhoades *et al.*, 2002). It was shown that plants expressing *ARF17* genes resistant to miR160 cleavage have increased levels of *ARF17* transcripts and altered levels of *GH3-like* mRNAs. Those plants also exhibited dramatic pleiotropic developmental defects such as leaf shape defects, premature inflorescence, reduced petal size, etc. Such phenotypes were also observed in plants expressing suppressors of RNA silencing or plants with mutations related to miRNA pathways (Mallory *et al.*, 2005). Mutant plants with miR160 resistant *ARF16* genes showed that miR160 and auxin independently regulate the activity of those genes, responsible for root cap development (Wang *et al.*, 2005). The transcription factor NAC1 is known to transduce auxin signals for lateral root emergence while also being a target of miR164. While measuring the levels of miR164 after auxin treatment, Guo *et al.* (2005) were able to detect a slight but consistent increase in the miRNA levels (*c.* 1.5-fold increase) some hours after treatment. This suggests a regulation of miR164 levels by auxin but also that the induction of miR164

by auxin may create a homeostatic mechanism that mediates clearance of *NAC1* mRNA after its initial induction by auxin. Overexpression of miR164 reduces lateral root formation, but an overexpression of miR164-resistant *NAC1* only slightly increases the number of lateral roots (Guo *et al.*, 2005). Furthermore, other groups reported that *NAC1* is a miR164 target but did not report a root phenotype when overexpressing miR164 or even fail to find evidence that miR164 targets *NAC1* *in vivo* (Laufs *et al.*, 2004; Mallory *et al.*, 2004a). The role of miR164 in the formation of lateral roots has to be cleared out.

MiR393 is predicted to target several F-box transcripts involved in the ubiquitination pathway (Bonnet *et al.*, 2004a; Jones-Rhoades & Bartel, 2004). Among those targets, cleavage products were found in *Arabidopsis* for *TRANSPORT INHIBITOR RESPONSE1 (TIR1)*; Jones-Rhoades & Bartel, 2004). This gene plays a central role in the auxin response pathway. TIR1 binds to AUX/IAA proteins, leading to an increased ubiquitination of the TIR1/AUX/IAA complex. In turn, this process will enhance the degradation of this complex, and release ARF proteins from repression, allowing auxin-responsive transcription (for review see Woodward & Bartel, 2005). It was shown recently that TIR1 is an auxin receptor that mediates AUX/IAA degradation and auxin-regulated transcription (Dharmasi *et al.*, 2005; Kepinski & Leyser, 2005). The discovery of an auxin receptor is a true landmark in the search for the mechanism of auxin action and the fact that this receptor is also a miRNA target highlights the importance of miRNAs as key regulators (Napier, 2005).

Regulation of the miRNA pathway Elevated levels of *DCL1* mRNAs were found in *dcl1* mutants, miRNA defective *ben1* mutants, and in plants expressing a virus-encoded suppressor of RNA silencing (P1/HC-PRO). Cleavage products corresponding to the activity of miR162 on *DCL1* transcripts were found, revealing a negative feedback regulation of this enzyme by a miRNA (Xie *et al.*, 2003). Transgenic plants expressing a mutated (but functional) *AGO1* mRNA with impaired complementarity to miR168 accumulated *AGO1* transcripts and showed developmental defects similar to those encountered in plants having crucial mutations for the miRNA pathway (*dcl1*, *gen1* or *hyl1*). Those defects could be rescued by the introduction of an artificial miRNA complementary to the mutated *AGO1* mRNA. These results demonstrate the existence of another feedback regulatory loop in the miRNA pathway (Vaucheret *et al.*, 2004).

Environmental and stress-related responses Despite an overwhelming propensity to target transcription factors, as previously mentioned, plant miRNAs are also predicted to match several other classes of targets. Some are linked to environmental changes or stress responses. For example, several ATP sulfurylase mRNAs (*APS1*, *APS3* or *APS4*) have a complementary site for miR395 (Bonnet *et al.*, 2004a;

Jones-Rhoades & Bartel, 2004). ATP sulfurylases catalyse the first step of inorganic sulfate assimilation. Products of degradation corresponding to the cleavage of *APS4* by miR395 were detected in *Arabidopsis*. It has been shown that miR395 is expressed upon sulfate starvation and that it is inversely correlated with *APS1* expression (Jones-Rhoades & Bartel, 2004). A very similar observation has recently been made regarding phosphate homeostasis, where it was shown that miR399 was induced by phosphate starvation, which in turn downregulated its target transcript encoding a ubiquitin-conjugating E2 enzyme through 5'UTR interaction (Chiou *et al.*, 2005; Fujii *et al.*, 2005). Accumulation of the *E2* transcripts was suppressed in transgenic plants overexpressing miR399. These transgenic plants accumulate inorganic phosphate and exhibit phosphate toxicity symptoms that phenocopy a loss-of-function *E2* mutant. This provides evidence that miR399 controls phosphate homeostasis by regulating a component of the proteolysis machinery in plants.

In poplar, a recent study revealed that the levels of many miRNAs cloned from woody tissues were either upregulated or downregulated in stem tissues submitted to mechanical stresses. This strongly suggests a role of miRNAs in tree defense systems against mechanical stresses (Lu *et al.*, 2005b). The miRNA mir397 is predicted to target laccases, a widespread family of enzymes conserved in bacteria, insects, plants and fungi (Bonnet *et al.*, 2004a; Jones-Rhoades & Bartel, 2004). Cleavage products for laccases were also found in *Arabidopsis* (Jones-Rhoades & Bartel, 2004). A homolog of miR397 was found in the recently released poplar genome (<http://genome.jgi-psf.org/Popptr1/Popptr1.home.html>) and is predicted to target 21 laccase homologs in this genome (E. Bonnet *et al.*, unpublished). This could be an interesting finding, as laccases are suspected to be involved in lignin (wood) formation (Mayer & Staples, 2002; Ranocha *et al.*, 2002).

Those results highlight what seems to be a typical feature of plant miRNAs described so far: a high degree of specificity, contrary to what is observed for animal miRNAs. A plant miRNA will typically regulate one or several members of a given protein family, usually closely related. This specificity was confirmed by expression experiments where the effect of the overexpression of a given plant miRNA was quantified using microarrays. Such experiments for five different miRNAs found a limited number mRNA differentially expressed when compared with controls where miRNAs were not overexpressed (Schwab *et al.*, 2005). Similar experiments in animals typically found a few hundred mRNAs differentially expressed per miRNA (Lim *et al.*, 2005). However, there is not always a one-to-one relationship. For example, the miRNAs miR319 and miR159 are grouped into one miRNA family, as they differ for only a few nucleotides. One regulates TCP transcription factors members (miR319) and the other MYB transcription factors (miR159) (Palatnik *et al.*, 2003). Another interesting aspect of miRNA regulation might be

their involvement in one or several major regulatory networks. For example, several miRNAs (and also ta-siRNAs: see next chapter) seem to play a key role in the auxin-signaling pathway (miR160, miR164, miR167 and miR393) and target different genes in this pathway (*ARFs*, *TIR1* and *NAC1*; see also Table 2). There is an indication for a combinatorial role for miRNAs here, in conjunction with other factors.

IV. Trans-acting siRNAs

In a screen for mutants impaired in the juvenile to adult phase transition, Peragine *et al.* (2004) identified several genes that were up-regulated in *sgs3*, *rdr6* and *ago7* mutants, including some AUXIN RESPONSE FACTORS (*ARF3* and *ARF4*). Among those they identified one locus that was silenced post-transcriptionally *in trans* by an endogenous siRNAs derived from a nonprotein-coding transcript. They also found that the process was SGS3-, RDR6- and DCL1-dependent, suggesting a relationship with the miRNA pathway.

An independent study by Vazquez *et al.* (2004b) was performed to identify the molecular basis of the *rdr6* and *sgs3* mutant phenotypes. A nonprotein-coding RNA transcript (now called *TAS1a*) was identified that accumulated in *rdr6* mutants. Small interfering RNAs were also identified that did not accumulate in *ago1*, *dcl1*, *ben1*, *hyl1*, *rdr6* and *sgs3* mutants. Vazquez *et al.* (2004b) showed that those siRNAs were processed from the *TAS1a* locus and that they guide the cleavage of several endogenous mRNAs.

Those two independent studies thus clearly established that siRNAs generated from noncoding transcripts were able to silence target mRNAs that have little overall resemblance to the gene from which they originate, demonstrating the existence of a third RNA silencing pathway, in addition to miRNAs and siRNAs, and providing yet another dimension to post-transcriptional mRNA regulation in plants (for a review see Vaucheret, 2005).

Later, it was shown that the cleavage of the noncoding transcript by a miRNA was a necessary step before the production of 21 nt siRNAs from the cleavage fragments (Allen *et al.*, 2005; Gascioli *et al.*, 2005; Yoshikawa *et al.*, 2005). There is also now evidence that DCL4 is the protein processing double-stranded noncoding ta-siRNA transcripts in 21-nt long ta-siRNAs (Gascioli *et al.*, 2005; Xie *et al.*, 2005b; Yoshikawa *et al.*, 2005).

A model for the processing of ta-siRNAs was proposed by some groups (Xie *et al.*, 2005b; Yoshikawa *et al.*, 2005): the miRNA cleaves a capped and polyadenylated transcript. Cleavage fragments (either 5' or 3') are bound by SGS3 or by proteins associated with SGS3, thus protecting them from degradation by enzymes acting on ssRNA. RDR6 then transforms the fragment into double stranded RNA that will be cleaved into 21 nt siRNAs by DCL4 (Fig. 2c).

The miRNA miR173 was found in wild-type *Arabidopsis*, but no experimental validation was done for its predicted

target, a protein of unknown function (Park *et al.*, 2002). miR390 was identified both by cloning and computational approaches (Bonnet *et al.*, 2004a; Sunkar & Zhu, 2004; Adai *et al.*, 2005) but predicted targets failed to be validated by 5' RACE experiments (Axtell & Bartel, 2005). More recently, using more complex target prediction algorithms and/or experimental approaches, the targets of miR173 and miR390 were identified as three noncoding transcript families encoding ta-siRNAs, designated *TAS1*, *TAS2* and *TAS3* (Peragine *et al.*, 2004; Vazquez *et al.*, 2004a; Allen *et al.*, 2005; Yoshikawa *et al.*, 2005).

The *TAS1* family is composed of three genes encoding a set of closely related ta-siRNAs that target four mRNAs of unknown function (Peragine *et al.*, 2004; Vazquez *et al.*, 2004a; Allen *et al.*, 2005). *TAS2*-derived ta-siRNAs target mRNAs encoding pentatricopeptide repeat (PPR) proteins (Allen *et al.*, 2005; Yoshikawa *et al.*, 2005). The *TAS3* locus specifies two ta-siRNAs that target a set of mRNAs corresponding to AUXIN RESPONSE FACTORS including *ARF3* and *ARF4* (Allen *et al.*, 2005; Williams *et al.*, 2005a). These loci are particularly interesting as Allen *et al.* (2005) found that miR390 genes, miR390 target sites, ta-siRNAs in *TAS3* primary transcripts and *TAS3* ta-siRNA target sites in *ARF3* and *ARF4* are conserved between several monocots and dicots, suggesting that this ta-siRNA pathway is at least 150 million or so years old.

Furthermore, other *ARF* genes are known to be regulated by miRNAs (Jones-Rhoades *et al.*, 2006; see also the above paragraph on miRNA targets), meaning that up to one third of the known *ARF* genes are regulated by miRNAs or ta-siRNAs. The association between auxin and small RNA regulation might suggest a need for a rapid clearance of auxin effectors mRNAs after signaling events (Bartel, 2004).

V. Conclusion and perspectives

It is clear that small RNAs hold many key functions in plants, particularly in genome stability, regulation of gene expression and defense. Some authors even compare them to the 'dark matter' of the universe because of their relatively recent discovery and their ubiquity. However, despite an impressive amount of knowledge acquired in a few years after their discovery, many aspects of plant small RNAs biogenesis and function remain unclear (Baulcombe, 2005; Carrington, 2005). Unlike many animals, plants encode multiple Dicer-like and RDR proteins. It was shown that this diversification contributed to specialization of small RNA-directed pathways. Nonetheless, the function of several key enzymes in those pathways remain unclear or unknown. For example, in *Arabidopsis*, seven out of 10 Argonaute family members (Fagard *et al.*, 2000; Carmell *et al.*, 2002) do show the characteristics of the proteins identified to be part of the miRNA (AGO1) or siRNA (AGO4) pathways (Liu *et al.*, 2004; Song *et al.*, 2004) and thus have the potential to form

alternative RISC complexes. In humans, four Argonaute proteins are equally competent to bind small RNAs, but only AGO2 is able to mediate cleavage. In *C. elegans*, the Argonaute family counts 23 members. Do they participate in some alternative small RNA pathways?

Very little is known about the ancient evolution of miRNAs. Allen *et al.* (2004) proposed an elegant model where miRNAs arise from inverted duplications of protein-coding genes. Here, the inverted duplications create a perfect hairpin secondary structure that is processed by Dicer enzymes into small RNAs, forming a set of siRNAs that will silence the gene from which they originate. Mutations will then occur in different parts of the hairpin-like structure, progressively transforming the siRNA into a miRNA, with its own locus and the possibility of silencing different targets. Alternative models are likely to exist to give rise to new microRNAs, but they remain to be discovered (Voinnet, 2004).

The inventory of miRNAs genes in different organisms is far from complete at the moment. One may expect the finding of new members in a range of species with specific environmental habits. Cloning and expression experiments together with computational analysis provide a complementary framework for the further identification of miRNAs. In this respect, the availability of newly annotated plant genome sequences will be an important resource for both experimental and *in silico* approaches. New and faster techniques for the deep sequencing of small RNAs in various organisms, next to analysis of their expression in specific arrays, will also help to get a complete picture of the small RNome.

The general view on miRNAs' autonomous function and interplay with other small RNAs within the plant cell is still fuzzy, despite some interesting hypotheses. For example, Bartel & Chen (2004) proposed that miRNAs could act as rheostats of gene expression. In plants, miRNAs could thus, for example, control redundant dose-sensitive genes following polyploidy events. This way, they could prevent the duplication of transcription factors from causing a hugely amplified response (Kidner & Martienssen, 2005). Another interesting hypothesis is that they could act as integrators of other genetic regulatory circuits rather than simple on-off switches. As more and more expression data become available, particularly from small RNA microarrays, it might be possible in the longer term to analyse and integrate those data in order to have an integrated view of the role of the different classes of small RNAs within the different cellular processes.

The question about the universality of the small RNAs and small RNA-driven processes among the eukaryotes and, as far as plants are concerned, among other members of the green lineage, needs further investigation. The same holds true for possible links with the transition of unicellular to multicellular organisms. Which small RNA-driven processes are absent from unicellular and colonial organisms? Did these never exist, or did they get lost?

The future obviously looks bright for biologists interested in plant small RNAs, with plenty of mysteries still to be unravelled.

Acknowledgements

We thank the anonymous referees for their constructive comments and apologize to those whose work was not included because of space constraints.

References

- Achard P, Herr A, Baulcombe DC, Harberd NP. 2004. Modulation of floral development by a gibberellin-regulated microRNA. *Development* **131**: 3357–3365.
- Adai A, Johnson C, Mlotshwa S, Archer-Evans S, Manocha V, Vance V, Sundaresan V. 2005. Computational prediction of miRNAs in *Arabidopsis thaliana*. *Genome Research* **15**: 78–91.
- Allen E, Xie Z, Gustafson AM, Sung GH, Spatafora JW, Carrington JC. 2004. Evolution of microRNA genes by inverted duplication of target gene sequences in *Arabidopsis thaliana*. *Nature Genetics* **36**: 1282–1290.
- Allen E, Xie Z, Gustafson AM, Carrington JC. 2005. microRNA-directed phasing during *trans*-acting siRNA biogenesis in plants. *Cell* **121**: 207–221.
- Ambros V, Bartel B, Bartel DP, Burge CB, Carrington JC, Chen X, Dreyfuss G, Eddy SR, Griffiths-Jones S, Marshall M, Matzke M, Ruvkun G, Tuschl T. 2003. A uniform system for microRNA annotation. *Rna* **9**: 277–279.
- Arazi T, Talmor-Neiman M, Stav R, Riese M, Huijser P, Baulcombe DC. 2005. Cloning and characterization of micro-RNAs from moss. *Plant Journal* **43**: 837–848.
- Argaman L, Hershberg R, Vogel J, Bejerano G, Wagner EG, Margalit H, Altuvia S. 2001. Novel small RNA-encoding genes in the intergenic regions of *Escherichia coli*. *Current Biology* **11**: 941–950.
- Aufsatz W, Mette MF, van der Winden J, Matzke M, Matzke AJ. 2002. HDA6, a putative histone deacetylase needed to enhance DNA methylation induced by double-stranded RNA. *EMBO Journal* **21**: 6832–6841.
- Aukerman MJ, Sakai H. 2003. Regulation of flowering time and floral organ identity by a MicroRNA and its APETALA2-like target genes. *Plant Cell* **15**: 2730–2741.
- Axtell MJ, Bartel DP. 2005. Antiquity of microRNAs and their targets in land plants. *Plant Cell* **17**: 1658–1673.
- Baker CC, Sieber P, Wellmer F, Meyerowitz EM. 2005. The early extra petals1 mutant uncovers a role for microRNA miR164c in regulating petal number in *Arabidopsis*. *Current Biology* **15**: 303–315.
- Bao N, Lye KW, Barton MK. 2004. MicroRNA binding sites in *Arabidopsis* class III HD-ZIP mRNAs are required for methylation of the template chromosome. *Developmental Cell* **7**: 653–662.
- Bartel DP. 2004. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* **116**: 281–297.
- Bartel DP, Chen CZ. 2004. Micromanagers of gene expression: the potentially widespread influence of metazoan microRNAs. *Nature Review of Genetics* **5**: 396–400.
- Baulcombe D. 2004. RNA silencing in plants. *Nature* **431**: 356–363.
- Baulcombe D. 2005. RNA silencing. *Trends in Biochemistry and Science* **30**: 290–293.
- Baumberger N, Baulcombe DC. 2005. *Arabidopsis* ARGONAUTE1 is an RNA Slicer that selectively recruits microRNAs and short interfering RNAs. *Proceedings of the National Academy of Sciences of the USA* **102**: 11928–11933.
- Bedell JA, Budiman MA, Nunberg A, Citek RW, Robbins D, Jones J, Flick

- E, Rholving T, Fries J, Bradford K, McMenamy J, Smith M, Holeman H, Roe BA, Wiley G, Korf IF, Rabinowicz PD, Lakey N, McCombie WR, Jeddeloh JA, Martienssen RA. 2005. Sorghum genome sequencing by methylation filtration. *PLoS Biology* 3: e13.
- Bentwich I, Avniel A, Karov Y, Aharonov R, Gilad S, Barad O, Barzilai A, Einat P, Einav U, Meiri E, Sharon E, Spector Y, Bentwich Z. 2005. Identification of hundreds of conserved and nonconserved human microRNAs. *Nature Genetics* 37: 766–770.
- Bernstein E, Caudy AA, Hammond SM, Hannon GJ. 2001. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* 409: 363–366.
- Billoud B, De Paeppe R, Baulcombe D, Boccara M. 2005. Identification of new small non-coding RNAs from tobacco and *Arabidopsis*. *Biochimie* 87: 905–910.
- Bohmer K, Camus I, Bellini C, Bouchez D, Caboche M, Benning C. 1998. AGO1 defines a novel locus of *Arabidopsis* controlling leaf development. *EMBO Journal* 17 (1): 170–180.
- Bollman KM, Aukerman MJ, Park MY, Hunter C, Berardini TZ, Poethig RS. 2003. HASTY, the *Arabidopsis* ortholog of exportin 5/MSN5, regulates phase change and morphogenesis. *Development* 130: 1493–1504.
- Bonnet E, Wuyts J, Rouzè P, Van de Peer Y. 2004a. Detection of 91 potential conserved plant microRNAs in *Arabidopsis thaliana* and *Oryza sativa* identifies important target genes. *Proceedings of the National Academy of Sciences, USA* 101: 11511–11516.
- Bonnet E, Wuyts J, Rouzè P, Van de Peer Y. 2004b. Evidence that microRNA precursors, unlike other non-coding RNAs, have lower folding free energies than random sequences. *Bioinformatics* 20: 2911–2917.
- Borsani O, Zhu J, Verslues PE, Sunkar R, Zhu JK. 2005. Endogenous siRNAs derived from a pair of natural cis-antisense transcripts regulate salt tolerance in *Arabidopsis*. *Cell* 123: 1279–1291.
- Boutet S, Vazquez F, Liu J, Béclin C, Fagard M, Gratias A, Morel JB, Créte P, Chen X, Vaucheret H. 2003. *Arabidopsis* HEN1: a genetic link between endogenous miRNA controlling development and siRNA controlling transgene silencing and virus resistance. *Current Biology* 13: 843–848.
- Britten RJ, Davidson EH. 1969. Gene regulation for higher cells: a theory. *Science* 165: 349–357.
- Carmell MA, Xuan Z, Zhang MQ, Hannon GJ. 2002. The Argonaute family: tentacles that reach into RNAi, developmental control, stem cell maintenance, and tumorigenesis. *Genes and Development* 16: 2733–2742.
- Carrington JC. 2005. Small RNAs and *Arabidopsis*. A fast forward look. *Plant Physiology* 138: 565–566.
- Chapman EJ, Prokhnovsky AI, Gopinath K, Dolja VV, Carrington JC. 2004. Viral RNA silencing suppressors inhibit the microRNA pathway at an intermediate step. *Genes and Development* 18: 1179–1186.
- Chen X. 2004. A microRNA as a translational repressor of APETALA2 in *Arabidopsis* flower development. *Science* 303: 2022–2025.
- Chiou TJ, Aung K, Lin SI, Wu CC, Chiang SF, Su CL. 2005. Regulation of phosphate homeostasis by microRNA in *Arabidopsis*. *Plant Cell* 18: 412–421.
- Clote P, Ferrè F, Kranakis E, Krizanc D. 2005. Structural RNA has lower folding energy than random RNA of the same dinucleotide frequency. *RNA* 11: 578–591.
- Cogoni C, Irelan JT, Schumacher M, Schmidhauser TJ, Selker EU, Macino G. 1996. Transgene silencing of the *al-1* gene in vegetative cells of *Neurospora* is mediated by a cytoplasmic effector and does not depend on DNA–DNA interactions or DNA methylation. *EMBO Journal* 15: 3153–3163.
- Couzin J. 2002. Breakthrough of the year. Small RNAs make big splash. *Science* 298: 2296–2297.
- Dalmay T, Hamilton A, Rudd S, Angell S, Baulcombe DC. 2000. An RNA-dependent RNA polymerase gene in *Arabidopsis* is required for posttranscriptional gene silencing mediated by a transgene but not by a virus. *Cell* 101: 543–553.
- Dharmasiri N, Dharmasiri S, Estelle M. 2005. The F-box protein TIR1 is an auxin receptor. *Nature* 435: 441–445.
- Dunoyer P, Voinnet O. 2005. The complex interplay between plant viruses and host RNA-silencing pathways. *Current Opinions in Plant Biology* 8: 415–423.
- Dunoyer P, Lecellier CH, Parizotto EA, Himber C, Voinnet O. 2004. Probing the microRNA and small interfering RNA pathways with virus-encoded suppressors of RNA silencing. *Plant Cell* 16: 1235–1250.
- Dunoyer P, Himber C, Voinnet O. 2005. DICER-LIKE 4 is required for RNA interference and produces the 21-nucleotide small interfering RNA component of the plant cell-to-cell silencing signal. *Nat Genetics* 37: 1356–1360.
- Dunoyer P, Himber C, Voinnet O. 2006. Induction, suppression and requirement of RNA silencing pathways in virulent *Agrobacterium tumefaciens* infections. *Nat Genetics* 38: 258–263.
- Eddy SR. 2001. Non-coding RNA genes and the modern RNA world. *Nature Review of Genetics* 2: 919–929.
- Elbashir SM, Harborth J, Weber K, Tuschl T. 2002. Analysis of gene function in somatic mammalian cells using small interfering RNAs. *Methods* 26: 199–213.
- Emery JF, Floyd SK, Alvarez J, Eshed Y, Hawker NP, Izhaki A, Baum SF, Bowman JL. 2003. Radial patterning of *Arabidopsis* shoots by class III HD-ZIP and KANADI genes. *Current Biology* 13: 1768–1774.
- Fagard M, Boutet S, Morel JB, Bellini C, Vaucheret H. 2000. AGO1, QDE-2, and RDE-1 are related proteins required for post-transcriptional gene silencing in plants, quelling in fungi, and RNA interference in animals. *Proceedings of the National Academy of Sciences, USA* 97: 11650–11654.
- Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE, Mello CC. 1998. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391: 806–811.
- Floyd SK, Bowman JL. 2004. Gene regulation: ancient microRNA target sequences in plants. *Nature* 428: 485–486.
- Friis EM, Pedersen KR, Crane PR. 2004. Araceae from the early Cretaceous of Portugal: evidence on the emergence of monocotyledons. *Proceedings of the National Academy of Sciences, USA* 101: 16565–16570.
- Fujii H, Chiou TJ, Lin SI, Aung K, Zhu JK. 2005. A miRNA involved in phosphate-starvation response in *Arabidopsis*. *Current Biology* 15: 2038–2043.
- Gascioli V, Mallory AC, Bartel DP, Vaucheret H. 2005. Partially redundant functions of *Arabidopsis* DICER-like enzymes and a role for DCL4 in producing trans-acting siRNAs. *Current Biology* 15: 1494–1500.
- Gendrel AV, Colot V. 2005. *Arabidopsis* epigenetics: when RNA meets chromatin. *Current Opinions in Plant Biology* 8: 142–147.
- Gesteland RF, Cech TR, Atkins JF. 1999. *The RNA world*. Cold Spring Harbor, NY, USA: Cold Spring Harbor Laboratory Press.
- Gilbert W. 1986. The RNA world. *Nature* 319: 618.
- Griffiths-Jones S, Grocock RJ, van Dongen S, Bateman A, Enright AJ. 2006. miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Research* 34: D140–D144.
- Guo S, Kemphues KJ. 1995. *par-1*, a gene required for establishing polarity in *C. elegans* embryos, encodes a putative Ser/Thr kinase that is asymmetrically distributed. *Cell* 81: 611–620.
- Guo HS, Xie Q, Fei JF, Chua NH. 2005. MicroRNA Directs mRNA cleavage of the transcription factor NAC1 to downregulate auxin signals for *Arabidopsis* lateral root development. *Plant Cell* 17: 1376–1386.
- Gustafson AM, Allen E, Givan S, Smith D, Carrington JC, Kasschau KD. 2005. ASRP: the *Arabidopsis* Small RNA Project Database. *Nucleic Acids Research* 33 (Database issue): D637–640.
- Hamilton AJ, Baulcombe DC. 1999. A species of small antisense RNA in posttranscriptional gene silencing in plants. *Science* 286: 950–952.
- Hamilton A, Voinnet O, Chappell L, Baulcombe D. 2002. Two classes of short interfering RNA in RNA silencing. *EMBO Journal* 21: 4671–4679.
- Hammond SM, Bernstein E, Beach D, Hannon GJ. 2000. An RNA-directed nuclease mediates post-transcriptional gene silencing in *Drosophila* cells. *Nature* 404: 293–296.
- Han MH, Goud S, Song L, Fedoroff N. 2004. The *Arabidopsis*

- double-stranded RNA-binding protein HYL1 plays a role in microRNA-mediated gene regulation. *Proceedings of the National Academy of Sciences, USA* 101: 1093–1098.
- Hannon GJ. 2002. RNA interference. *Nature* 418: 244–251.
- Herr AJ, Jensen MB, Dalmay T, Baulcombe DC. 2005. RNA polymerase IV directs silencing of endogenous DNA. *Science* 308: 118–120.
- Hiraguri A, Itoh R, Kondo N, Nomura Y, Aizawa D, Murai Y, Koiwa H, Seki M, Shinozaki K, Fukuhara T. 2005. Specific interactions between Dicer-like proteins and HYL1/DRB-family dsRNA-binding proteins in *Arabidopsis thaliana*. *Plant Molecular Biology* 57: 173–188.
- Hofacker IL, Fontana W, Stadler PF, Bonhoeffer LS, Tacker M, Schuster P. 1994. Fast Folding and Comparison of RNA Secondary Structures. *Monatshefte für Chemie* 125: 167–188.
- Huttenhofer A, Kieffmann M, Meier-Ewert S, O'Brien J, Lehrach H, Bachelier JP, Brosius J. 2001. RNomics: an experimental approach that identifies 201 candidates for novel, small, non-messenger RNAs in mouse. *EMBO Journal* 20: 2943–2953.
- Jackson JP, Lindroth AM, Cao X, Jacobsen SE. 2002. Control of CpNpG DNA methylation by the KRYPTONITE histone H3 methyltransferase. *Nature* 416: 556–560.
- Jacob F, Monod J. 1961. Genetic regulatory mechanisms in the synthesis of proteins. *Journal of Molecular Biology* 3: 318–356.
- Jones-Rhoades MW, Bartel DP. 2004. Computational identification of plant microRNAs and their targets, including a stress-induced miRNA. *Molecular Cell* 14: 787–799.
- Jones-Rhoades MW, Bartel DP, Bartel B. 2006. MicroRNAs and Their Regulatory Roles in Plants. *Annual Review of Plant Biology* 57: 19–53.
- Juarez MT, Kui JS, Thomas J, Heller BA, Timmermans MC. 2004. microRNA-mediated repression of rolled leaf1 specifies maize leaf polarity. *Nature* 428: 84–88.
- Kanno T, Huettel B, Mette MF, Aufsatz W, Jaligot E, Daxinger L, Kreil DP, Matzke M, Matzke AJ. 2005. Atypical RNA polymerase subunits required for RNA-directed DNA methylation. *Nat Genetics* 37: 761–765.
- Kasschau KD, Xie Z, Allen E, Llave C, Chapman EJ, Krizan KA, Carrington JC. 2003. P1/HC-Pro, a viral suppressor of RNA silencing, interferes with *Arabidopsis* development and miRNA function. *Developmental Cell* 4: 205–217.
- Kepinski S, Leyser O. 2005. The *Arabidopsis* F-box protein TIR1 is an Auxin receptor. *Nature* 435: 446–451.
- Kidner CA, Martienssen RA. 2004. Spatially restricted microRNA directs leaf polarity through ARGONAUTE1. *Nature* 428: 81–84.
- Kidner CA, Martienssen RA. 2005. The developmental role of microRNA in plants. *Current Opinions in Plant Biology* 8: 38–44.
- Kim J, Jung JH, Reyes JL, Kim YS, Kim SY, Chung KS, Kim JA, Lee M, Lee Y, Narry Kim V, Chua NH, Park CM. 2005. microRNA-directed cleavage of ATHB15 mRNA regulates vascular development in *Arabidopsis* inflorescence stems. *Plant Journal* 42: 84–94.
- van der Krol AR, Mur LA, Beld M, Mol JN, Stuitje AR. 1990. Flavonoid genes in petunia: addition of a limited number of gene copies may lead to a suppression of gene expression. *Plant Cell* 2: 291–299.
- Kurihara Y, Takashi Y, Watanabe Y. 2006. The interaction between DCL1 and HYL1 is important for efficient and precise processing of pri-miRNA in plant microRNA biogenesis. *Rna* 12: 206–212.
- Kurihara Y, Watanabe Y. 2004. *Arabidopsis* micro-RNA biogenesis through Dicer-like 1 protein functions. *Proceedings of the National Academy of Sciences, USA* 101: 12753–12758.
- Lagos-Quintana M, Rauhut R, Lendeckel W, Tuschl T. 2001. Identification of novel genes coding for small expressed RNAs. *Science* 294: 853–858.
- Lakatos L, Szittyá G, Silhavy D, Burgyan J. 2004. Molecular mechanism of RNA silencing suppression mediated by p19 protein of tombusviruses. *EMBO Journal* 23: 876–884.
- Lau NC, Lim LP, Weinstein EG, Bartel DP. 2001. An abundant class of tiny RNAs with probable regulatory roles in *Caenorhabditis elegans*. *Science* 294: 858–862.
- Laufs P, Peaucelle A, Morin H, Traas J. 2004. MicroRNA regulation of the CUC genes is required for boundary size control in *Arabidopsis* meristems. *Development* 131: 4311–4322.
- Lauter N, Kampani A, Carlson S, Goebel M, Moose SP. 2005. microRNA172 down-regulates glossy15 to promote vegetative phase change in maize. *Proceedings of the National Academy of Sciences, USA* 102: 9412–9417.
- Lee RC, Ambros V. 2001. An extensive class of small RNAs in *Caenorhabditis elegans*. *Science* 294: 862–864.
- Lee RC, Feinbaum RL, Ambros V. 1993. The *C. elegans* heterochronic gene lin-4 encodes small RNAs with antisense complementarity to lin-14. *Cell* 75: 843–854.
- Lee R, Feinbaum R, Ambros V. 2004a. A short history of a short RNA. *Cell* 116: S89–92. following S96.
- Li H, Xu L, Wang H, Yuan Z, Cao X, Yang Z, Zhang D, Xu Y, Huang H. 2005. The putative RNA-dependent RNA polymerase RDR6 acts synergistically with ASYMMETRIC LEAVES1 and 2 to repress BREVIPEDICELLUS and MicroRNA165/166 in *Arabidopsis* leaf development. *Plant Cell* 17: 2157–2171.
- Lim LP, Glasner ME, Yekta S, Burge CB, Bartel DP. 2003. Vertebrate microRNA genes. *Science* 299: 1540.
- Lim LP, Lau NC, Garrett-Engle P, Grimson A, Schelter JM, Castle J, Bartel DP, Linsley PS, Johnson JM. 2005. Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs. *Nature* 433: 769–773.
- Lippman Z, Martienssen R. 2004. The role of RNA interference in heterochromatic silencing. *Nature* 431: 364–370.
- Lippman Z, May B, Yordan C, Singer T, Martienssen R. 2003. Distinct mechanisms determine transposon inheritance and methylation via small interfering RNA and histone modification. *PLoS Biology* 1: E67.
- Liu J, Carmell MA, Rivas FV, Marsden CG, Thomson JM, Song JJ, Hammond SM, Joshua-Tor L, Hannon GJ. 2004. Argonaute2 is the catalytic engine of mammalian RNAi. *Science* 305: 1437–1441.
- Llave C, Kasschau KD, Rector MA, Carrington JC. 2002a. Endogenous and silencing-associated small RNAs in plants. *Plant Cell* 14: 1605–1619.
- Llave C, Xie Z, Kasschau KD, Carrington JC. 2002b. Cleavage of Scarecrow-like mRNA targets directed by a class of *Arabidopsis* miRNA. *Science* 297: 2053–2056.
- Lu C, Fedoroff N. 2000. A mutation in the *Arabidopsis* HYL1 gene encoding a dsRNA binding protein affects responses to abscisic acid, Auxin, and cytokinin. *Plant Cell* 12: 2351–2366.
- Lu C, Tej SS, Luo S, Haudenschild CD, Meyers BC, Green PJ. 2005a. Elucidation of the small RNA component of the transcriptome. *Science* 309: 1567–1569.
- Lu S, Sun YH, Shi R, Clark C, Li L, Chiang VL. 2005b. Novel and mechanical stress-responsive microRNAs in *Populus trichocarpa* that are absent from *Arabidopsis*. *Plant Cell* 17: 2186–2203.
- Mallory AC, Reinhart BJ, Bartel D, Vance VB, Bowman LH. 2002. A viral suppressor of RNA silencing differentially regulates the accumulation of short interfering RNAs and micro-RNAs in tobacco. *Proceedings of the National Academy of Sciences, USA* 99: 15228–15233.
- Mallory AC, Dugas DV, Bartel DP, Bartel B. 2004a. MicroRNA regulation of NAC-domain targets is required for proper formation and separation of adjacent embryonic, vegetative, and floral organs. *Current Biology* 14: 1035–1046.
- Mallory AC, Reinhart BJ, Jones-Rhoades MW, Tang G, Zamore PD, Barton MK, Bartel DP. 2004b. MicroRNA control of PHABULOSA in leaf development: importance of pairing to the microRNA 5' region. *EMBO Journal* 23: 3356–3364.
- Mallory AC, Bartel DP, Bartel B. 2005. MicroRNA-directed regulation of *Arabidopsis* AUXIN RESPONSE FACTOR17 is essential for proper development and modulates expression of early auxin response genes. *Plant Cell* 17: 1360–1375.
- Mayer AM, Staples RC. 2002. Laccase: new functions for an old enzyme. *Phytochemistry* 60: 551–565.

- McHale NA, Koning RE. 2004. MicroRNA-directed cleavage of *Nicotiana sylvestris* PHAVOLUTA mRNA regulates the vascular cambium and structure of apical meristems. *Plant Cell* 16: 1730–1740.
- Mette MF, van der Winden J, Matzke M, Matzke AJ. 2002. Short RNAs can identify new candidate transposable element families in *Arabidopsis*. *Plant Physiology* 130: 6–9.
- Millar AA, Gubler F. 2005. The *Arabidopsis* GAMYB-like genes, MYB33 and MYB65, are MicroRNA-regulated genes that redundantly facilitate anther development. *Plant Cell* 17: 705–721.
- Montgomery MK, Xu S, Fire A. 1998. RNA as a target of double-stranded RNA-mediated genetic interference in *Caenorhabditis elegans*. *Proceedings of the National Academy of Sciences, USA* 95: 15502–15507.
- Morel JB, Godon C, Mourrain P, Béclin C, Boutet S, Feuerbach F, Proux F, Vaucheret H. 2002. Fertile hypomorphic ARGONAUTE (ago1) mutants impaired in post-transcriptional gene silencing and virus resistance. *Plant Cell* 14: 629–639.
- Mourrain P, Béclin C, Elmayan T, Feuerbach F, Godon C, Morel JB, Jouette D, Lacombe AM, Nikic S, Picault N, Remoue K, Sanial M, Vo TA, Vaucheret H. 2000. *Arabidopsis* SGS2 and SGS3 genes are required for posttranscriptional gene silencing and natural virus resistance. *Cell* 101: 533–542.
- Napier RM. 2005. TIRs of joy: new receptors for auxin. *Bioessays* 27: 1213–1217.
- Napoli C, Lemieux C, Jorgensen R. 1990. Introduction of a chimeric chalcone synthase gene into petunia results in reversible co-suppression of homologous genes *in trans*. *Plant Cell* 2: 279–289.
- Nykanen A, Haley B, Zamore PD. 2001. ATP requirements and small interfering RNA structure in the RNA interference pathway. *Cell* 107: 309–321.
- Olivas WM, Muhlrud D, Parker R. 1997. Analysis of the yeast genome: identification of new non-coding and small ORF-containing RNAs. *Nucleic Acids Research* 25: 4619–4625.
- Onodera Y, Haag JR, Ream T, Nunes PC, Pontes O, Pikaard CS. 2005. Plant nuclear RNA polymerase IV mediates siRNA and DNA methylation-dependent heterochromatin formation. *Cell* 120: 613–622.
- Palatnik JF, Allen E, Wu X, Schommer C, Schwab R, Carrington JC, Weigel D. 2003. Control of leaf morphogenesis by microRNAs. *Nature* 425: 257–263.
- Palauqui JC, Elmayan T, Pollien JM, Vaucheret H. 1997. Systemic acquired silencing: transgene-specific post-transcriptional silencing is transmitted by grafting from silenced stocks to non-silenced scions. *EMBO Journal* 16: 4738–4745.
- Papp I, Mette MF, Aufsatz W, Daxinger L, Schauer SE, Ray A, van der Winden J, Matzke M, Matzke AJ. 2003. Evidence for nuclear processing of plant micro RNA and short interfering RNA precursors. *Plant Physiology* 132: 1382–1390.
- Parizotto EA, Dunoyer P, Rahm N, Himber C, Voinnet O. 2004. *In vivo* investigation of the transcription, processing, endonucleolytic activity, and functional relevance of the spatial distribution of a plant miRNA. *Genes and Development* 18: 2237–2242.
- Park MY, Wu G, Gonzalez-Sulser A, Vaucheret H, Poethig RS. 2005. Nuclear processing and export of microRNAs in *Arabidopsis*. *Proceedings of the National Academy of Sciences of the USA* 102: 3691–3696.
- Park W, Li J, Song R, Messing J, Chen X. 2002. CARPEL FACTORY, a Dicer homolog, and HEN1, a novel protein, act in microRNA metabolism in *Arabidopsis thaliana*. *Current Biology* 12: 1484–1495.
- Pasquinelli AE, Reinhart BJ, Slack FJ, Martindale MQ, Kuroda MI, Maller B, Hayward DC, Ball EE, Degnan B, Muller P, Spring J, Srinivasan A, Fishman M, Finnerty J, Corbo J, Levine M, Leahy P, Davidson E, Ruvkun G. 2000. Conservation of the sequence and temporal expression of *let-7* heterochronic regulatory RNA. *Nature* 408: 86–89.
- Peragine A, Yoshikawa M, Wu G, Albrecht HL, Poethig RS. 2004. SGS3 and SGS2/SDE1/RDR6 are required for juvenile development and the production of *trans*-acting siRNAs in *Arabidopsis*. *Genes and Development* 18: 2368–2379.
- Pham JW, Pellino JL, Lee YS, Carthew RW, Sontheimer EJ. 2004. A Dicer-2-dependent 80s complex cleaves targeted mRNAs during RNAi in *Drosophila*. *Cell* 117: 83–94.
- Plasterk RH. 2002. RNA silencing: the genome's immune system. *Science* 296: 1263–1265.
- Pontier D, Yahubyan G, Vega D, Bulski A, Saez-Vasquez J, Hakimi MA, Lerbs-Mache S, Colot V, Lagrange T. 2005. Reinforcement of silencing at transposons and highly repeated sequences requires the concerted action of two distinct RNA polymerases IV in *Arabidopsis*. *Genes and Development* 19: 2030–2040.
- Ranocha P, Chabannes M, Chamayou S, Danoun S, Jauneau A, Boudet AM, Goffner D. 2002. Laccase down-regulation causes alterations in phenolic metabolism and cell wall structure in poplar. *Plant Physiology* 129: 145–155.
- Reinhart BJ, Slack FJ, Basson M, Pasquinelli AE, Bettinger JC, Rougvie AE, Horvitz HR, Ruvkun G. 2000. The 21-nucleotide *let-7* RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature* 403: 901–906.
- Reinhart BJ, Weinstein EG, Rhoades MW, Bartel B, Bartel DP. 2002. MicroRNAs in plants. *Genes and Development* 16: 1616–1626.
- Rhoades MW, Reinhart BJ, Lim LP, Burge CB, Bartel B, Bartel DP. 2002. Prediction of plant microRNA targets. *Cell* 110: 513–520.
- Rivas E, Klein RJ, Jones TA, Eddy SR. 2001. Computational identification of noncoding RNAs in *E. coli* by comparative genomics. *Current Biology* 11: 1369–1373.
- Ruvkun G, Wightman B, Ha I. 2004. The 20 years it took to recognize the importance of tiny RNAs. *Cell* 116: S93–6. following S96.
- Schauer SE, Jacobsen SE, Meinke DW, Ray A. 2002. DICER-LIKE1: blind men and elephants in *Arabidopsis* development. *Trends in Plant Science* 7: 487–491.
- Schwab R, Palatnik JF, Riester M, Schommer C, Schmid M, Weigel D. 2005. Specific effects of microRNAs on the plant transcriptome. *Developmental Cell* 8: 517–527.
- Schwarz DS, Du Hutvagner GT, Xu Z, Aronin N, Zamore PD. 2003. Asymmetry in the assembly of the RNAi enzyme complex. *Cell* 115: 199–208.
- Simon-Mateo C, Garcia JA. 2006. MicroRNA-guided processing impairs Plum pox virus replication, but the virus readily evolves to escape this silencing mechanism. *Journal of Virology* 80: 2429–2436.
- Slack FJ, Basson M, Liu Z, Ambros V, Horvitz HR, Ruvkun G. 2000. The *lin-41* RBCC gene acts in the *C. elegans* heterochronic pathway between the *let-7* regulatory RNA and the LIN-29 transcription factor. *Molecular Cell* 5: 659–669.
- Smith CJ, Watson CF, Bird CR, Ray J, Schuch W, Grierson D. 1990. Expression of a truncated tomato polygalacturonase gene inhibits expression of the endogenous gene in transgenic plants. *Molecular General Genetics* 224: 477–481.
- Song JJ, Smith SK, Hannon GJ, Joshua-Tor L. 2004. Crystal structure of Argonaute and its implications for RISC slicer activity. *Science* 305: 1434–1437.
- Soutschek J, Akinc A, Bramlage B, Charisse K, Constien R, Donoghue M, Elbashir S, Geick A, Hadwiger P, Harborth J, John M, Kesavan V, Lavine G, Pandey RK, Racie T, Rajeev KG, Rohl I, Toudjarska I, Wang G, Wuschko S, Bumcrot D, Kotliansky V, Limmer S, Manoharan M, Vornlocher HP. 2004. Therapeutic silencing of an endogenous gene by systemic administration of modified siRNAs. *Nature* 432: 173–178.
- Sunkar R, Zhu JK. 2004. Novel and stress-regulated microRNAs and other small RNAs from *Arabidopsis*. *Plant Cell* 16: 2001–2019.
- Sunkar R, Girke T, Jain PK, Zhu JK. 2005. Cloning and characterization of MicroRNAs from rice. *Plant Cell* 17: 1397–1411.
- Tang G, Reinhart BJ, Bartel DP, Zamore PD. 2003. A biochemical framework for RNA silencing in plants. *Genes and Development* 17: 49–63.
- Tomari Y, Matranga C, Haley B, Martinez N, Zamore PD. 2004. A protein sensor for siRNA asymmetry. *Science* 306: 1377–1380.

- Tuskan GA, DiFazio SP, Teichmann T. 2004. Poplar genomics is getting popular: the impact of the poplar genome project on tree research. *Plant Biology (Stuttgart)* 6: 2–4.
- Vaucheret H. 2005. MicroRNA-dependent *trans*-acting siRNA production. *Sci STKE* 2005: pe43.
- Vaucheret H, Vazquez F, Cr  t   P, Bartel DP. 2004. The action of ARGONAUTE1 in the miRNA pathway and its regulation by the miRNA pathway are crucial for plant development. *Genes and Development* 18: 1187–1197.
- Vaughn MW, Martienssen RA. 2005. Finding the right template: RNA Pol IV, a plant-specific RNA polymerase. *Molecular Cell* 17: 754–756.
- Vazquez F, Gasciolli V, Cr  t   P, Vaucheret H. 2004a. The nuclear dsRNA binding protein HYL1 is required for microRNA accumulation and plant development, but not posttranscriptional transgene silencing. *Current Biology* 14: 346–351.
- Vazquez F, Vaucheret H, Rajagopalan R, Lepers C, Gasciolli V, Mallory AC, Hilbert JL, Bartel DP, Cr  t   P. 2004b. Endogenous *trans*-acting siRNAs regulate the accumulation of *Arabidopsis* mRNAs. *Molecular Cell* 16: 69–79.
- Voynet O. 2004. Shaping small RNAs in plants by gene duplication. *Nature Genetics* 36: 1245–1246.
- Voynet O. 2005. Non-cell autonomous RNA silencing. *FEBS Letters* 579: 5858–5871.
- Voynet O, Baulcombe DC. 1997. Systemic signalling in gene silencing. *Nature* 389: 553.
- Voynet O, Pinto YM, Baulcombe DC. 1999. Suppression of gene silencing: a general strategy used by diverse DNA and RNA viruses of plants. *Proceedings of the National Academy of Sciences, USA* 96: 14147–14152.
- Wang MB, Bian XY, Wu LM, Liu LX, Smith NA, Isenegger D, Wu RM, Masuta C, Vance VB, Watson JM, Rezaian A, Dennis ES, Waterhouse PM. 2004b. On the role of RNA silencing in the pathogenicity and evolution of viroids and viral satellites. *Proceedings of the National Academy of Sciences, USA* 101: 3275–3280.
- Wang XJ, Reyes JL, Chua NH, Gaasterland T. 2004a. Prediction and identification of *Arabidopsis thaliana* microRNAs and their mRNA targets. *Genome Biology* 5: R65.
- Wang JW, Wang LJ, Mao YB, Cai WJ, Xue HW, Chen XY. 2005. Control of root cap formation by MicroRNA-targeted Auxin response factors in *Arabidopsis*. *Plant Cell* 2005: 17 (8): 2204–2216.
- Wassarman KM, Repoila F, Rosenow C, Storz G, Gottesman S. 2001. Identification of novel small RNAs using comparative genomics and microarrays. *Genes and Development* 15: 1637–1651.
- Wassenegger M, Heimes S, Riedel L, Sanger HL. 1994. RNA-directed *de novo* methylation of genomic sequences in plants. *Cell* 76: 567–576.
- Wightman B, Ha I, Ruvkun G. 1993. Posttranscriptional regulation of the heterochronic gene *lin-14* by *lin-4* mediates temporal pattern formation in *C. elegans*. *Cell* 75: 855–862.
- Williams L, Carles CC, Osmont KS, Fletcher JC. 2005a. A database analysis method identifies an endogenous *trans*-acting short-interfering RNA that targets the *Arabidopsis* ARF2, ARF3, and ARF4 genes. *Proceedings of the National Academy of Sciences, USA* 102: 9703–9708.
- Williams L, Grigg SP, Xie M, Christensen S, Fletcher JC. 2005b. Regulation of *Arabidopsis* shoot apical meristem and lateral organ formation by microRNA miR166g and its AtHD-ZIP target genes. *Development* 132: 3657–3668.
- Woodward AW, Bartel B. 2005. A receptor for auxin. *Plant Cell* 17: 2425–2429.
- Wu-Scharf D, Jeong B, Zhang C, Cerutti H. 2000. Transgene and transposon silencing in *Chlamydomonas reinhardtii* by a DEAH-box RNA helicase. *Science* 290: 1159–1162.
- Xie Z, Fan B, Chen C, Chen Z. 2001. An important role of an inducible RNA-dependent RNA polymerase in plant antiviral defense. *Proceedings of the National Academy of Sciences, USA* 98: 6516–6521.
- Xie Z, Kasschau KD, Carrington JC. 2003. Negative feedback regulation of *Dicer-Like 1* in *Arabidopsis* by microRNA-guided mRNA degradation. *Current Biology* 13: 784–789.
- Xie Z, Johansen LK, Gustafson AM, Kasschau KD, Lellis AD, Zilberman D, Jacobsen SE, Carrington JC. 2004. Genetic and functional diversification of small RNA pathways in plants. *PLoS Biology* 2: E104.
- Xie Z, Allen E, Fahlgren N, Calamar A, Givan SA, Carrington JC. 2005a. Expression of *Arabidopsis* MIRNA genes. *Plant Physiology* 138: 2145–2154.
- Xie Z, Allen E, Wilken A, Carrington JC. 2005b. DICER-LIKE 4 functions in *trans*-acting small interfering RNA biogenesis and vegetative phase change in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* 102: 12984–12989.
- Ye K, Malinina L, Patel DJ. 2003. Recognition of small interfering RNA by a viral suppressor of RNA silencing. *Nature* 426: 874–878.
- Yu B, Yang Z, Li J, Minakhina S, Yang M, Padgett RW, Steward R, Chen X. 2005. Methylation as a crucial step in plant microRNA biogenesis. *Science* 307: 932–935.
- Yoo BC, Kragler F, Varkonyi-Gasic E, Haywood V, Archer-Evans S, Lee YM, Lough TJ, Lucas WJ. 2004. A systemic small RNA signaling system in plants. *Plant Cell* 16: 1979–2000.
- Yoshikawa M, Peragine A, Park MY, Poethig RS. 2005. A pathway for the biogenesis of *trans*-acting siRNAs in *Arabidopsis*. *Genes and Development* 19: 2164–2175.
- Zamore PD. 2002. Ancient pathways programmed by small RNAs. *Science* 296: 1265–1269.
- Zhong R, Ye ZH. 2004. Amphivasal vascular bundle 1, a gain-of-function mutation of the IFL1/REV gene, is associated with alterations in the polarity of leaves, stems and carpels. *Plant Cell Physiology* 45: 369–385.
- Zilberman D, Cao X, Jacobsen SE. 2003. ARGONAUTE4 control of locus-specific siRNA accumulation and DNA and histone methylation. *Science* 299: 716–719.
- Zilberman D, Cao X, Johansen LK, Xie Z, Carrington JC, Jacobsen SE. 2004. Role of *Arabidopsis* ARGONAUTE4 in RNA-directed DNA methylation triggered by inverted repeats. *Current Biology* 14: 1214–1220.
- Zuker M, Stiegler P. 1981. Optimal computer folding of large RNA sequences using thermodynamics and auxiliary information. *Nucleic Acids Research* 9: 133–148.



About *New Phytologist*

- *New Phytologist* is owned by a non-profit-making **charitable trust** dedicated to the promotion of plant science, facilitating projects from symposia to open access for our Tansley reviews. Complete information is available at www.newphytologist.org.
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as-ready' via *OnlineEarly* – the 2004 average submission to decision time was just 30 days. Online-only colour is **free**, and essential print colour costs will be met if necessary. We also provide 25 offprints as well as a PDF for each article.
- For online summaries and ToC alerts, go to the website and click on 'Journal online'. You can take out a **personal subscription** to the journal for a fraction of the institutional price. Rates start at £109 in Europe/\$202 in the USA & Canada for the online edition (click on 'Subscribe' at the website).
- If you have any questions, do get in touch with Central Office (newphytol@lancaster.ac.uk; tel +44 1524 594691) or, for a local contact in North America, the US Office (newphytol@ornl.gov; tel +1 865 576 5261).